

PHARMACOKINETICS, EFFICACY AND SAFETY OF MIRIKIZUMAB AS INDUCTION THERAPY IN PAEDIATRIC PATIENTS WITH MODERATELY TO SEVERELY ACTIVE ULCERATIVE COLITIS: RESULTS FROM THE PHASE 2 SHINE-1 STUDY

Amy Larkin

UEG Week Copenhagen 2023

Poster

IBD

October 15, 2023

Phase 2 SHINE-1 study reports the first efficacy and safety data for mirikizumab, a p19 IL-23 inhibitor, in 26 pediatric patients aged 2 to <18 years with moderately to severely active ulcerative colitis after 12 weeks of induction therapy.

KEY POINTS

- At Week 12, MMS clinical response was achieved in 69.2% of pediatric patients, clinical remission in 38.5%, and endoscopic remission in 53.8%, with PUCAI response in 76.9% and PUCAI remission in 38.5%.
- These efficacy rates were consistent with or numerically higher than those observed in adults treated with 300 mg mirikizumab in the Phase 3 LUCENT-1 study (adult MMS clinical response 63.5%, clinical remission 25.6%, endoscopic remission 36.3%).
- Pharmacokinetic analysis confirmed that dosing of 300 mg IV for patients >40 kg and 5 mg/kg IV for patients ≤40 kg achieved exposures similar to adult levels, while 10 mg/kg yielded approximately 2-fold higher exposure.
- Safety findings in pediatric patients were consistent with the acceptable safety profile observed in the adult LUCENT-1 induction study.
- This material is relevant for pediatric gastroenterologists and researchers interested in novel biologics for pediatric inflammatory bowel disease and the development pathway for mirikizumab in children with ulcerative colitis.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/7c78fb78-743c-11ee-83ac-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.