

ENDOSCOPIC ULTRASONOGRAPHY IS A GAME CHANGER IN PANCREATIC CYSTS MANAGEMENT: INTENTION-TO-TREAT ANALYSIS

Andrea Caravati

UEG Week Copenhagen 2023

Poster

Pancreas

October 15, 2023

Endoscopic ultrasonography changed presumptive diagnosis in 36.9% and treatment indication in 28.5% of 355 patients with presumed pancreatic cysts at a single center between 2019 and 2021.

KEY POINTS

- EUS identified pancreatic ductal adenocarcinoma in 5.6% of patients in the cohort.
- Diagnosis change was most frequent for rare benign or non-neoplastic cysts (52.6%), IPMN with HRS (47.2%), and cNET (46.7%).
- All radiologically presumed solid pseudopapillary tumors were confirmed by EUS, with no change in diagnosis or treatment.
- All presumed serous cystic neoplasms in which EUS changed diagnosis had an atypical macrocystic multilocular presentation.
- The authors conclude that the impact of EUS is sufficiently remarkable that performing this exam should be considered at least once during evaluation of pancreatic cysts.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/26be09bc-743c-11ee-a175-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.