

CORRELATION OF SIMPLE LABORATORY MARKERS WITH ENDOSCOPIC REMISSION IN PATIENTS WITH CROHN'S DISEASE

Andreas Protopapas

UEG Week Copenhagen 2023

Poster

IBD

October 15, 2023

A study of 76 treated Crohn's disease patients found that mean platelet volume and neutrophil-to-lymphocyte ratio correlated significantly with endoscopic disease activity measured by CDEIS, suggesting potential value as non-invasive markers of endoscopic remission.

KEY POINTS

- In 76 patients (mean age 38.4 years) who underwent colonoscopy and laboratory testing within three months, endoscopic remission was present in 44 patients by CDEIS and 36 by SES-CD
- Mean platelet volume showed statistically significant correlation with both CDEIS-based endoscopic stage ($p=0.01$) and presence of endoscopic remission ($p=0.008$)
- Neutrophil-to-lymphocyte ratio correlated significantly with CDEIS-based endoscopic stage ($p=0.022$) and remission status ($p=0.037$), while plateletcrit correlated with endoscopic stage ($p=0.036$)
- Endoscopic remission status based on SES-CD did not show statistically significant correlation with any of the laboratory indices tested
- The authors conclude that further investigation of NLR and MPV in larger patient cohorts is warranted

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/23003d00-743b-11ee-a082-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.