

EPPOEM: A RETROSPECTIVE MULTICENTER COMPARATIVE ANALYSIS OF SELECTIVE INNER MUSCLE LAYER MYOTOMY VERSUS FULL-THICKNESS MYOTOMY IN THE ENDOSCOPIC TREATMENT OF ACHALASIA USING POEM PROCEDURE

Arthur berger

UEG Week Copenhagen 2023

Poster

Oesophagus

October 15, 2023

A retrospective bi-centric study of 158 achalasia patients found no significant difference in clinical efficacy or GERD outcomes between selective inner layer myotomy and full-thickness myotomy during POEM, but full-thickness myotomy resulted in more postoperative pain and longer hospitalization.

KEY POINTS

- Clinical efficacy at 6 months was similar between selective inner layer myotomy and full-thickness myotomy groups (84% versus 90%, $p=0.57$), as was efficacy at 12 months (70% versus 80%, $p=0.74$)
- Full-thickness myotomy patients required more opioid analgesics than selective inner layer myotomy patients (45% versus 21%, $p<0.01$) and had longer hospital stays (2.94 versus 2.17 days, $p<0.001$)
- Rates of esophagitis at 6 months were comparable between groups (16% selective inner layer versus 12% full-thickness, $p=0.73$), with no significant difference in heartburn at 6 or 12 months
- The authors conclude that selective internal myotomy should be preferred over full-thickness myotomy based on the similar efficacy but improved postoperative profile

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/d303f7d6-743c-11ee-8455-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.