

FEASIBILITY, SAFETY AND EFFICACY OF ENDOSCOPIC SUBMUCOSAL DISSECTION FOR RESIDUAL OR RECURRENT SUPERFICIAL RECTAL NEOPLASTIC LESIONS AFTER TRANSANAL SURGERY.

Arthur berger

UEG Week Copenhagen 2023

Poster

Colorectal

October 15, 2023

A multicenter retrospective study of 38 patients found that endoscopic submucosal dissection achieved 92% en bloc resection and 68% curative resection rates for recurrent or incomplete rectal neoplasms after transanal surgery, with only one residual lesion at follow-up.

KEY POINTS

- En bloc resection was achieved in 92% of cases (35/38), with R0 and curative resection rates both at 68% (26/38), despite 78% of lesions showing severe fibrosis from prior surgery.
- At median 6.2 months follow-up, only one residual lesion (3%, 1/38) was detected, which was a low-grade dysplasia adenoma from one of the three non-en bloc resections.
- Complications included four post-operative bleeding events and two stenoses; eleven patients had rectal fat visualized during resection but none required surgery, with median hospital stay of 1.8 days.
- Median lesion size was 44 mm, median procedure time was 79 minutes, and pathology revealed 32 adenomas and 3 T2 adenocarcinomas among the 38 resected lesions.
- This retrospective study may interest endoscopists managing recurrent rectal neoplasms after transanal surgical approaches.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/f565a474-743b-11ee-94b6-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.