

IMPACT OF GLUTEN-FREE DIET ON PSYCHOLOGICAL ISSUES ASSOCIATED WITH CELIAC DISEASE

Asfold Parfenov

UEG Week Copenhagen 2023

Poster

Small Intestine & Nutrition

October 15, 2023

A cross-sectional study of 48 celiac disease patients found high rates of anxiety (72.8% with anxiety reactions, 43.7% at clinically significant levels) but no association between anxiety and adherence to a gluten-free diet.

KEY POINTS

- Patients with celiac disease showed decreased physical and mental quality of life components compared to 20 controls without gastrointestinal disease (36.1 and 40.0 respectively, $p < 0.01$).
- Anxiety was more prevalent than depression in celiac disease patients, with 72.8% experiencing anxiety reactions and 43.7% reaching clinically significant anxiety levels on the Hospital Anxiety and Depression Scale.
- No significant differences in frequency or severity of anxiety were found between patients adhering to a gluten-free diet and those not adhering.
- Depression symptoms were more frequent in patients following a gluten-free diet, though not reaching clinically significant levels, and the severity of depression increased in this group.
- This study is relevant for clinicians managing the psychological aspects of celiac disease and understanding the complex relationship between dietary adherence and mental health outcomes.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/199c55a8-743d-11ee-8194-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.