

PANCREATIC DUCTAL FLUID AND HCO₃⁻ SECRETION INCREASE IN DIABETES IN WHICH CFTR PLAYS A CENTRAL ROLE

Attila Ébert

UEG Week Copenhagen 2023

Poster

Pancreas

October 15, 2023

A mouse model study found that diabetes increases both basal and stimulated pancreatic ductal fluid and bicarbonate secretion, contrary to some prior reports, with CFTR and other ion transporters playing a central role.

KEY POINTS

- Basal fluid and bicarbonate secretion were significantly elevated in diabetic mice compared to normal mice.
- Secretin-stimulated fluid secretion was also significantly higher in diabetic mice.
- Inhibition of CFTR significantly reduced bicarbonate secretion in both normal and diabetic mice.
- Expression of ion and water transporters including CFTR, ANO1, NHE-1, and AQP1 increased in diabetes.
- The study used streptozotocin-induced diabetes in wild type and CFTR knockout mice with measurements by fluorescence microscopy, real-time PCR, and immunohistochemistry.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/09f9a5b6-743c-11ee-a8c4-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.