

INTEGRATED RELAXATION PRESSURE ON HIGH RESOLUTION MANOMETRY DIFFERS BETWEEN MEASUREMENT ACROSS LOWER ESOPHAGEAL SPHINCTER ALONE VS. CRURAL DIAPHRAGM IN THE PRESENCE OF A HIATUS HERNIA

C Prakash Gyawali

UEG Week Copenhagen 2023

Poster

Oesophagus

October 15, 2023

Differential measurement of integrated relaxation pressure across the lower esophageal sphincter and crural diaphragm separately reduces over-diagnosis of esophagogastric junction outflow obstruction in patients with hiatus hernia.

KEY POINTS

- In 467 patients undergoing high resolution manometry, when hiatus hernia was present, IRP measured across both LES and CD together was significantly higher than IRP measured separately across LES or CD ($p < 0.001$).
- Of 11 patients with IRP greater than 15 mmHg in type 2 and type 3 EGJ using standard methodology, only 3 retained persistent IRP elevation when LES and CD were assessed separately.
- Standard methodology found similar median IRP values across the three EGJ morphological subtypes: type 1 (no hernia) 6.1 mmHg, type 2 (LES-CD separation < 3 cm) 5.5 mmHg, and type 3 (separation ≥ 3 cm) 8.5 mmHg.
- The authors speculate that standard IRP measurement may capture pressurization within the hiatus hernia itself when measured across both LES and CD in hernia patients.
- This study is relevant for clinicians interpreting high resolution manometry in patients with hiatus hernia to avoid misclassifying patients as having outflow obstruction.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/9b8306e4-743c-11ee-acc5-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.