gutflix
GI learning, personalised

Summary

ROME IV-BASED CHRONIC NAUSEA AND VOMITING SYNDROME AND FUNCTIONAL DYSPEPSIA FREQUENTLY OVERLAP AMONG PATIENTS WITH INVESTIGATED DYSPEPSIA

Chloé Melchior

UEG Week Copenhagen 2023 | Poster

Oesophagus

October 17, 2023

In investigated dyspepsia, 73.2% of chronic nausea and vomiting syndrome patients also meet functional dyspepsia criteria, and those with overlap exhibit worse quality of life, higher depression, and worse sleep than either condition alone.

KEY POINTS

- Among 2,317 patients with investigated dyspepsia, 651 had functional dyspepsia alone, 144 had chronic nausea and vomiting syndrome alone, and 393 met criteria for both disorders.
- 73.2% of patients with chronic nausea and vomiting syndrome fulfilled criteria for functional dyspepsia, while 37.6% of functional dyspepsia patients met criteria for chronic nausea and vomiting syndrome.
- Patients with overlap of both conditions had significantly worse quality of life, higher depression levels, and worse sleep quality than patients with either diagnosis alone.
- All symptoms were at least as prevalent in overlap patients as in functional dyspepsia alone, except early satiation.

VIEW THIS CONTENT ON GUTFLIX

https://gutflix.eu/search/d/b7c8ec1a-743c-11ee-8206-0242ac140004

AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.