

COMMUNICATION GAP BETWEEN PATIENTS AND HEALTH CARE PROFESSIONALS ON BOWEL URGENCY WITH FOCUS ON CROHN'S DISEASE: RESULTS FROM THE COMMUNICATING NEEDS AND FEATURES OF IBD EXPERIENCES (CONFIDE) SURVEY

Simon Travis

UEG Week Copenhagen 2023

Poster

IBD

October 15, 2023

The CONFIDE survey reveals that bowel urgency is the second most common symptom among patients with moderate-to-severe Crohn's disease but is underrecognized by healthcare professionals, indicating a significant communication gap.

KEY POINTS

- Bowel urgency was reported by 42% of US and 38% of EU5 patients with Crohn's disease as a current symptom, making it the second most common symptom after diarrhea.
- Among patients receiving advanced therapies, 46% of US and 39% of EU5 patients still experienced bowel urgency, yet only 19% of US and 16% of EU5 healthcare professionals ranked it among the top three symptoms.
- Most patients who experienced bowel urgency were not completely comfortable discussing it with healthcare professionals (only 40% US and 27% EU5), with embarrassment being the primary barrier (73% US, 64% EU5).
- One-third of US and 40% of EU5 healthcare professionals reported not proactively discussing bowel urgency, most commonly because they expected patients to raise the issue themselves.
- This analysis of surveys from 215 US and 547 EU5 patients and 200 US and 503 EU5 healthcare professionals highlights a communication gap that may lead to underappreciation of bowel urgency's impact on patients' daily lives.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/a7683aba-743c-11ee-8278-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.