

THE EFFECTS OF BIFIDOBACTERIUM BREVE, BIF195 ON NON-STEROIDAL ANTI-INFLAMMATORY DRUG-INDUCED GASTROINTESTINAL INJURY – THE INFLUENCE OF DOSE

Eamonn Quigley

UEG Week Copenhagen 2023

Poster

Oesophagus

October 15, 2023

A randomized controlled trial found that a low dose of the probiotic *Bifidobacterium breve* Bif195 (15 billion CFU/day) did not reduce ibuprofen-induced small intestinal mucosal injury compared to placebo in 160 healthy adults.

KEY POINTS

- Lewis score (measure of small intestinal damage) showed no statistically significant difference between Bif195 and placebo groups after 6 weeks of ibuprofen 800 mg/day ($p=0.474$).
- Ibuprofen induced markedly lower mucosal damage than aspirin in previous studies (Lewis score AUC 2208-2358 vs 4351), which may have limited the trial's ability to detect a protective effect.
- Two prior trials showed that higher doses of Bif195 (50-100 billion CFU/day) alleviated aspirin-induced mucosal injury in the small intestine and stomach.
- Authors suggest the results may indicate a dose-related effect but caution that differences in NSAIDs and trial design limit direct comparison across the three studies.
- The probiotic was well-tolerated with no treatment-related adverse events; study assessed 160 healthy adults (63% women, mean age 27 years) using video capsule endoscopy.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/7ae3ad9a-743b-11ee-97bd-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.