

THE USE OF ALGINATE DURING PROTON PUMP INHIBITOR WASHOUT IMPROVES PPI DEPRESCRIBING BY IMPROVING SYMPTOM BURDEN AND QUALITY OF LIFE

Edoardo Vincenzo Savarino

UEG Week Copenhagen 2023

Poster

Oesophagus

October 15, 2023

A retrospective study of 620 patients on long-term PPI therapy found that alginate use during mandatory PPI washout before reflux testing enabled 74% of patients with typical symptoms and 54% with atypical symptoms to discontinue PPIs at follow-up.

KEY POINTS

- Patients were asked to stop PPIs for at least 2 weeks prior to reflux testing and allowed to take alginate as rescue therapy, with 80% using alginate during withdrawal.
- At follow-up (median 2-3 months after testing), 280 of 380 patients with typical symptoms and 130 of 240 patients with atypical symptoms did not resume PPIs.
- Symptom scores improved significantly after testing, with GerdQ-positive patients decreasing from 64% to 48% for typical symptoms and RSI-positive patients decreasing from 72% to 61% for atypical symptoms.
- Patients taking alginate on a standardised schedule (after each meal and before night) were able to stop PPIs more frequently than those using it as needed (76% vs 48%).
- The study is relevant to clinicians managing patients on long-term PPI therapy and considering deprescribing strategies; the lead author has financial relationships with the alginate manufacturer.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/f417a8e2-743b-11ee-9c6f-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.