

CIRCULATING MIRNAS IN INFLAMMATORY BOWEL DISEASES - PRELIMINARY RESULTS

Elena Milanesi

UEG Week Copenhagen 2023

Poster

IBD

October 15, 2023

Preliminary screening of 179 plasma microRNAs in 6 IBD patients versus 12 controls identified 45 differentially expressed microRNAs, including 29 upregulated and 7 downregulated, with let-7a-5p and miR-30b-5p correlating with disease duration.

KEY POINTS

- 45 microRNAs were differentially expressed ($p < 0.05$) in the plasma of IBD patients compared to non-IBD controls, with 29 showing overexpression (fold regulation > 1.5) and 7 showing underexpression (fold regulation < -1.5).
- let-7a-5p was upregulated in the IBD group (fold regulation 2.45, $p = 0.001$) and positively correlated with years of disease duration ($p = 0.01$, Pearson $r = 0.919$).
- miR-30b-5p also showed positive correlation with years of disease duration ($p = 0.049$, Pearson $r = 0.812$).
- The authors state these are preliminary results from a small cohort (6 IBD, 12 controls) that require validation in a larger IBD cohort to assess potential value as diagnostic and prognostic biomarkers.
- This material is most relevant to researchers investigating microRNA biomarkers in inflammatory bowel disease and those interested in early-phase biomarker discovery studies.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/7b872308-743b-11ee-9a91-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.