

14-DAY TAILORED PCR-GUIDED TRIPLE THERAPY VERSUS 14-DAY NON-BISMUTH CONCOMITANT QUADRUPLE THERAPY FOR HELICOBACTER PYLORI ERADICATION: A MULTICENTER, OPEN-LABEL RANDOMIZED CONTROLLED TRIAL, THE HEPYSE TRIAL

Frédéric HELUWAERT

UEG Week Copenhagen 2023

Poster

Stomach & H. Pylori

October 15, 2023

A French multicentre randomised controlled trial demonstrated non-inferiority of 14-day PCR-guided tailored triple therapy compared to 14-day non-Bismuth concomitant quadruple therapy for first-line H. pylori eradication in a high clarithromycin resistance setting.

KEY POINTS

- In the intention-to-treat population of 241 randomised patients, eradication rates were 99.2% for tailored PCR-guided therapy (esomeprazole, amoxicillin, and clarithromycin or levofloxacin based on resistance) versus 95.9% for standard concomitant quadruple therapy (esomeprazole, amoxicillin, clarithromycin, and metronidazole), with a difference of +3.30% and lower CI bound of -0.68%.
- Per-protocol analysis showed eradication rates of 98.8% versus 96.5%, with difference of +2.35% and lower CI bound of -3.33%, both meeting the prespecified non-inferiority margin of -5%.
- The tailored PCR-guided approach was associated with significantly fewer adverse events (41.2% versus 61.4%, $p=0.003$), particularly gastrointestinal adverse events (36.0% versus 57.9%, $p=0.001$).
- This study may inform clinicians treating H. pylori infection in regions with high clarithromycin resistance who have access to rapid susceptibility testing.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/8a9f4ee6-743c-11ee-b0fb-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.