

EFFECT OF PROBIOTIC (VIVOMIXX) ON GUT MICROBIOME AND SHORT CHAIN FATTY ACIDS IN HEALTHY ASIAN VOLUNTEERS - A PILOT EXPLORATORY RANDOMIZED CONTROLLED TRIAL

Fria May Manejero

UEG Week Copenhagen 2023

Poster

Colorectal

October 15, 2023

A pilot randomized controlled trial in 31 healthy Asian adults found that the probiotic vivomixx altered gut microbiome composition, with rifaximin pre-treatment enhancing short-chain fatty acid production while colonic lavage pre-treatment deterred probiotic bacterial engraftment.

KEY POINTS

- All three interventions—polyethylene glycol colonic lavage, rifaximin, and vivomixx probiotic—resulted in large microbiome composition changes.
- Pre-treatment with colonic lavage deterred engraftment, shown by smaller increases in Bifidobacterium abundance compared to vivomixx alone.
- Rifaximin pre-treatment showed selective engraftment of Lactobacillus and increased concentrations of acetic, butyric, and propionic acids, sustained at 8 weeks.
- Vivomixx demonstrated favorable impact on gut microbiome composition but minimal impact on short-chain fatty acid concentrations in this study.
- The authors suggest these findings provide insights for future trial design and that gut microbiome and short-chain fatty acid modulation is a potential treatment target in dysbiosis-associated diseases.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/0f3fbbb4-743c-11ee-8643-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.