gutflix
GI learning, personalised

Summary

PREVALENCE OF SEXUAL DYSFUNCTION IN PATIENTS WITH LIVER DISEASE

Hicham El Bacha

UEG Week Vienna 2024 | Poster

Hepatobiliary

October 13, 2024

A 2023 monocentric study of 120 male patients found significantly higher rates of sexual dysfunction across all domains in those with chronic liver disease compared to controls, with cirrhosis identified as the only statistically significant risk factor.

KEY POINTS

- Male patients with chronic liver disease (n=60) had significantly higher rates of erectile dysfunction (66.6% vs 26.6%, p=0.003), orgasmic dysfunction (58.3% vs 23.3%, p=0.012), disorders of desire (68.3% vs 16.6%, p<0.001), disorders of sexual satisfaction (63.3% vs 30%, p<0.001), and disorders of overall satisfaction (75% vs 28.3%, p<0.001) compared to controls without liver pathology (n=60).
- Cirrhosis was the only statistically significant risk factor for sexual dysfunction identified in the study (OR 7.9, p=0.012), and 65% of the liver disease cohort were cirrhotics.
- All patients (100%) reported never having been asked about their sexual health during a consultation.
- The authors conclude that sexual dysfunction should be systematically screened in liver disease patients with a multidisciplinary approach to management.

VIEW THIS CONTENT ON GUTFLIX

https://gutflix.eu/search/d/714cf09e-9112-11ef-8a0f-0242ac140004

AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.