

NOVEL MAGNIFYING ENDOSCOPIC FINDING, CAPILLARIES PUSHED UP BY CLOUDY MUCOSA (CPCM) MAY BE SUPPORTIVE MARKER FOR POST-HELICOBACTER PYLORI EARLY GASTRIC CANCER

Hideki Kobara

UEG Week Copenhagen 2023

Poster

Oesophagus

October 15, 2023

A retrospective study of 238 gastric lesions found that scattered capillaries pushed up by cloudy mucosa (CPCM) detected on magnifying endoscopy with narrow band imaging showed higher specificity and positive predictive value for diagnosing early gastric cancer after *Helicobacter pylori* eradication compared to HP-infected early gastric cancer.

KEY POINTS

- Among 103 cancers and 63 non-cancers in HP eradicated lesions, CPCM specificity was 98.5% and positive predictive value was 98.2%, compared to 83.3% and 84.8% respectively in HP-infected lesions.
- CPCM was defined as three or more cloudy mucosa in one maximum magnification ME-NBI image and was present in 33.7% of HP eradicated lesions versus 45.8% of HP-infected lesions.
- The difference in specificity between groups was statistically significant ($p=0.0126$), as was the difference in positive predictive value ($p=0.0248$).
- The authors suggest CPCM may serve as a supportive diagnostic marker for HP eradicated early gastric cancer, which is difficult to diagnose due to low atypical epithelium mimicking non-cancerous structure.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflifx.eu/search/d/20292682-743b-11ee-9776-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.