

ENDOFLIP TAILORED SELECTIVE CIRCULAR POEM REDUCES SHORT TERM POST-PROCEDURAL GASTRO-ESOPHAGEAL REFLUX DISEASE SYMPTOMS IN TREATMENT NAÏVE PATIENTS BUT NOT IN PREVIOUSLY TREATED

Homira Ayubi

UEG Week Copenhagen 2023

Poster

Oesophagus

October 15, 2023

In 96 achalasia patients undergoing POEM, short-term GERD developed in 18.7% overall, with previous treatment independently predicting GERD (OR 7.4) rather than whether the procedure was EndoFLIP-tailored.

KEY POINTS

- GERD occurred in 18.7% of patients at 3 months post-POEM and did not differ significantly between EndoFLIP-tailored (21.1%) and non-tailored (17.2%) procedures.
- Previous treatment was the only independent predictor of GERD on multivariate analysis (OR 7.412, $p=0.004$), with rates of 26.8% in treated versus 12.7% in treatment-naïve patients.
- Among tailored-POEM patients, previously treated patients had significantly higher GERD rates than naïve patients (42.9% versus 8.3%, $p=0.034$), whereas no such difference existed in the non-tailored group.
- Higher baseline distensibility index correlated with GERD development in tailored-procedure patients (2.9 ± 1.7 versus 2.0 ± 1.5 , $p=0.038$), but myotomy length and approach (anterior versus posterior) did not correlate with GERD.
- This study is relevant to endoscopists performing POEM who wish to understand GERD risk stratification, particularly in patients with prior achalasia treatments.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/279d1a94-743c-11ee-92c0-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.