

# THIOPURINES IN CROHN'S DISEASE: DO THEY STILL HAVE A PLACE AS MONOTHERAPY IN REAL LIFE?

Ikram Errabih

UEG Week Copenhagen 2023

Poster

IBD

October 15, 2023

**A retrospective study of 407 Crohn's disease patients on thiopurines found that while monotherapy achieved clinical remission in 87% of cases, combination with biotherapy was the only independent predictor of mucosal healing and showed superior CRP normalization.**

## KEY POINTS

- Among 375 patients on thiopurine monotherapy and 32 on combination therapy, clinical remission rates were similar (86.93% versus 75%,  $p=0.08$ ), but combination therapy was an independent predictor of mucosal healing in multivariate analysis ( $OR=1.8$ ,  $p=0.01$ ).
- CRP normalization was significantly higher with combination therapy (84.37%) compared to monotherapy (39.73%,  $p<0.001$ ), while fecal calprotectin negativity was lower with combination therapy (13.63% versus 41.86%,  $p=0.015$ ).
- Thiopurines were indicated as maintenance treatment in 66.33% of cases, as monotherapy for fistulizing forms when biotherapy was unavailable (before 2016) in 25.8%, and for cortico-dependency in 7.86%.
- Metabolite monitoring was rarely performed, and when done (43 cases), 6-TGN was in therapeutic range in only 32.55% of cases, requiring therapeutic adjustment.
- This retrospective analysis from 2012 to 2022 may interest clinicians evaluating the role of thiopurine monotherapy versus combination therapy in contemporary Crohn's disease management.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/1497f9d6-743d-11ee-a487-0242ac140004>



#### **AI-generated summary**

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.