

COLON CANCER IN DIFFERENT ETHNIC GROUPS, POPULATION-BASED LARGE COHORT OF TWO ETHNIC GROUPS IN ISRAEL

Jabir Alkrenawi

UEG Week Copenhagen 2023

Poster

Colorectal

October 15, 2023

A retrospective study of 61,679 colorectal cancer patients in Israel found that Arab patients were diagnosed significantly younger than Jewish patients, with 21.3% diagnosed at age 50 or younger compared to 8% of Jewish patients.

KEY POINTS

- Arab CRC patients had a mean diagnosis age of 62.5 years versus 70.8 years for Jewish patients, with higher rates of diabetes (37.6% vs 27.5%) and obesity (36% vs 21.4%).
- All-cause mortality was lower among Arab patients (49.6% vs 60.8%), though they died at a younger mean age (70.9 vs 79.8 years).
- Arab patients had lower rates of ischemic heart disease (9.7% vs 12.6%), hypertension (45.6% vs 53.7%), and chronic kidney disease (8.1% vs 10.3%).
- The authors conclude that disparities in risk factors, comorbidities, and mortality between ethnic groups warrant special consideration for specific populations in CRC screening and management.
- This material is relevant for gastroenterologists and oncologists working with ethnically diverse populations or interested in health disparities in colorectal cancer.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflif.eu/search/d/0281dc4a-743c-11ee-9814-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.