

LONG TERM SINGLE CENTER COMPREHENSIVE ANALYSIS OF POST-POEM GASTROESOPHAGEAL REFLUX AND ITS SEQUELAE IN MORE THAN 500 PATIENTS

Jan Martinek

UEG Week Copenhagen 2023

Poster

Oesophagus

October 15, 2023

A retrospective analysis of 519 patients who underwent peroral endoscopic myotomy (POEM) for achalasia shows that reflux esophagitis rates decline from 39.1% at 3 months to 20.5% at 6 years, but approximately half of patients require long-term proton pump inhibitor therapy.

KEY POINTS

- Among 390 patients tested at 3 months post-POEM, 45.6% had abnormal acid exposure on 24-hour pH monitoring off proton pump inhibitors.
- Reflux esophagitis decreased over time: 39.1% at 3 months, 35.7% at 24-36 months, and 20.5% at 6 years, with severe esophagitis (LA grade C/D) declining from 4.3% at 3 months to 0% at 6 years.
- At 48, 60, and 72 months, approximately 53%, 48%, and 54% of patients were taking proton pump inhibitors, with 80% using them daily.
- Severe reflux complications were rare: no new cases of Barrett's esophagus or esophageal adenocarcinoma occurred, and one patient developed a peptic stricture that was successfully treated with balloon dilation.
- The authors conclude that patients after POEM should remain under long-term surveillance despite the low risk of reflux complications.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/75b9e720-743c-11ee-a3a5-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.