

ISOLATED CECAL NECROSIS – A RARE DIFFERENTIAL DIAGNOSIS OF A COMMON CLINICAL PRESENTATION

José Damasceno

UEG Week Copenhagen 2023

Poster

Colorectal

October 15, 2023

A 51-year-old man with diabetes and end-stage kidney disease on hemodialysis presented with right lower quadrant pain and diarrhea, diagnosed with isolated cecal necrosis confirmed by colonoscopy and biopsy, successfully managed conservatively.

KEY POINTS

- The patient presented with 12-hour history of right lower quadrant abdominal pain and bloodless diarrhea, with leukocytosis 12200/ul and C-reactive protein 190 mg/L.
- Abdominal CT scan showed 10 cm diffuse parietal thickening of the ascending colon with mesenteric fat densification, suggesting inflammatory pathology.
- Colonoscopy revealed black-colored cecum mucosa and ileocecal valve with superficial ulceration and abrupt transition to normal mucosa; histology confirmed severe ischemic colitis with necrosis.
- Conservative management with fluid therapy and antibiotics was chosen given absence of CT complications and patient surgical risk, resulting in favorable clinical evolution.
- Isolated cecal necrosis is a rare variant of ischemic colitis, more common in elderly patients with comorbidities, and represents an uncommon cause of acute abdomen with right lower quadrant pain.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/04a900ec-743d-11ee-aaf3-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.