

COMBINATION OF GRANULOCYTE-MONOCYTE APHERESIS AND TOFACITINIB: MULTICENTRE AND RETROSPECTIVE STUDY

Jose Luis Cabriada Nuño

UEG Week Copenhagen 2023

Poster

IBD

October 15, 2023

A retrospective multicentre study of 12 patients with ulcerative colitis found that combining granulocyte-monocyte apheresis with tofacitinib after primary non-response or loss of response was associated with clinical improvement and no adverse events.

KEY POINTS

- Partial Mayo score significantly decreased one month after the last GMA session ($p=0.027$) in patients who had not responded to tofacitinib alone.
- All patients receiving steroids at baseline were able to discontinue them, and no patient required colectomy during follow-up.
- Patients received a median of 11 GMA sessions (IQR 3-20), with 50% continuing maintenance GMA, and 73% started GMA after primary non-response to tofacitinib.
- No adverse events related to the combination therapy were reported, though 36% of patients were switched to a new therapy during follow-up.
- This small retrospective study may inform clinicians managing refractory ulcerative colitis patients with prior exposure to multiple biologics.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflīx.eu/search/d/28fc6df0-743b-11ee-b44f-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.