

EFFICACY OF INTRAVENOUS FERRIC CARBOXYMALTOSE TREATMENT IS INDEPENDENT FROM DISEASE ACTIVITY AND DOES NOT REQUIRE THERAPY OPTIMISATION IN CASE OF SILENT IBD WITH IDA – A PROSPECTIVE STUDY

Klaudia Farkas

UEG Week Copenhagen 2023

Poster

IBD

October 15, 2023

Prospective study of 73 IBD patients with iron deficiency anaemia found that intravenous ferric-carboxymaltose increased haemoglobin levels equally in active disease and remission groups without requiring treatment modification in patients in remission.

KEY POINTS

- 73 IBD patients (46 Crohn's disease, 16 ulcerative colitis, mean age 40.4 years) with iron deficiency anaemia were divided into active disease with therapy modification (n=27), remission with iron alone (n=23), and remission with therapy optimisation (n=23) groups
- Haemoglobin levels increased significantly in all groups ($p<0.001$) at weeks 8, 16, and 24, with no difference between active and remission groups or between the two remission subgroups
- Ferritin, serum iron levels, and transferrin saturation increased in all groups ($p<0.05$) without differences between groups, and no adverse events were observed
- The authors conclude that intravenous ferric-carboxymaltose is effective in remission without treatment modification, though they recommend treatment modification should still be considered due to the pathomechanism of iron deficiency anaemia in IBD

VIEW THIS CONTENT ON GUTFLIX

<https://gutflifx.eu/search/d/0e1b7f7a-743c-11ee-9b5a-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.