

ISOLATED ANASTOMOTIC LINE ULCERATIONS ARE ASSOCIATED WITH A HIGHER RISK OF POSTOPERATIVE RECURRENCE IN PATIENTS WITH CROHN'S DISEASE UNDERGOING ILEOCECAL RESECTION AND A DIFFERENTIAL MUCOSA-ASSOCIATED MICROBIOME COMPOSITION

Krzysztof Borowski

UEG Week Copenhagen 2023

Poster

IBD

October 15, 2023

In Crohn's disease patients with otherwise healthy neo-terminal ileum after ileocolonic resection, isolated anastomotic ulcers were associated with a five-fold increased risk of postoperative recurrence compared to patients without ulceration.

KEY POINTS

- Among 60 Crohn's disease patients undergoing ileocolonic resection with no ulcerations in the neo-terminal ileum at first postoperative colonoscopy, those with ulcerations limited to the anastomotic line (modified Rutgeerts score i2a, 43.3% of patients) had significantly higher risk of postoperative recurrence than those without any ulceration (adjusted HR 5.15, 95% CI 2.24-11.87, $p=0.00012$) over a median 4.2-year follow-up.
- Postoperative recurrence was defined as a composite outcome including clinical recurrence with objective inflammation, change in CD-related therapy, endoscopic balloon dilation, or intestinal resection.
- Microbiome analysis showed differential bacterial composition on both sides of the anastomosis between groups, with *Klebsiella*, *Fusobacterium*, and *Lachnospira* more abundant in patients with anastomotic ulcers, while *Lachnospiraceae* NK4A136 was more abundant in patients without ulceration.
- This prospective multicenter study provides evidence for gastroenterologists managing postoperative Crohn's disease patients regarding the prognostic significance of isolated anastomotic ulcers.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/794f9ea2-743c-11ee-a3fc-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.