

COMPARISON OF THE RESULTS OF MAGNETICALLY CONTROLLED CAPSULE ENDOSCOPY AND HIGH-DEFINITION GASTROSCOPY WITH PROPOFOL SEDATION: A PROSPECTIVE STUDY IN PATIENTS WITH UNINVESTIGATED FUNCTIONAL DYSPESIA

László Madácsy

UEG Week Copenhagen 2023

Poster

Oesophagus

October 15, 2023

A prospective study of 89 patients with functional dyspepsia found that magnetic-controlled capsule endoscopy (MCCE) and conventional gastroscopy showed comparable diagnostic accuracy for gastric lesions, with MCCE demonstrating superior patient tolerability.

KEY POINTS

- MCCE and gastroscopy showed no significant differences in detecting polyps, ulcers, submucosal tumours, early gastric cancer, lymphoma, or pangastritis in 89 patients (mean age 45) examined between 2018 and 2023.
- MCCE was more accurate than gastroscopy for detecting minimal foveolar hyperplasia (9 vs 2, $p=0.01$) and proximal gastritis (4 vs 1, $p=0.02$), while gastroscopy detected more antral gastritis cases (66 vs 53, $p=0.03$).
- All patients reported MCCE was more tolerable than gastroscopy with propofol sedation based on questionnaire assessment of discomfort, pain, and anxiety levels.
- The authors conclude MCCE may be a useful alternative to gastroscopy in patients under age 50 with uninvestigated functional dyspepsia to help select those requiring biopsy and potentially reduce endoscopy waiting lists.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/8c62fade-743c-11ee-8bff-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.