

THE ROLE AND PREVALENCE OF CANDIDA IN GUT MICROBIOME IN PATIENTS WITH NAFLD

Lidiia Mulka

UEG Week Copenhagen 2023

Poster

Hepatobiliary

October 15, 2023

A case-control study of 93 NAFLD patients and 85 controls found Candida prevalence was nearly three times higher in NAFLD patients (35% vs 13%) and correlated with elevated triglycerides, TNF-alpha, and an increased Firmicutes/Bacteroidetes ratio.

KEY POINTS

- Candida prevalence in the gut microbiota of NAFLD patients was 35% compared to 13% in controls without fatty liver infiltration
- NAFLD patients showed a higher Firmicutes/Bacteroidetes ratio (4.02 vs 1.82) and lower Bacteroidetes levels (21.10 vs 45.54) compared to controls
- In NAFLD patients, Candida presence positively correlated with the Firmicutes/Bacteroidetes index ($r=0.63$), triglycerides ($r=0.56$), and TNF-alpha ($r=0.8$), while no such correlations were observed in controls
- The Firmicutes/Bacteroidetes ratio in NAFLD patients correlated with increases in triglycerides ($r=0.53$), ALT ($r=0.61$), and VLDL ($r=0.4$)
- The study used quantitative real-time PCR to characterize gut microbiome composition in patients diagnosed with NAFLD by liver transient elastography

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/31bf5838-743d-11ee-8911-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.