

IMPACT OF SMOKING HABIT ON THE OUTCOMES OF ENDOSCOPIC SLEEVE GASTROPLASTY

Maria Valeria Matteo

UEG Week Copenhagen 2023

Poster

Small Intestine & Nutrition

October 15, 2023

A retrospective analysis of 275 endoscopic sleeve gastroplasty patients over 24 months found that smoking status (smoker, non-smoker, or ex-smoker) did not significantly affect weight loss outcomes or safety.

KEY POINTS

- Weight loss metrics (total body weight loss, excess weight loss, and absolute weight loss) were similar across smokers, non-smokers, and ex-smokers at 6, 12, and 24 months following endoscopic sleeve gastroplasty.
- No major perioperative complications were reported in any of the three smoking status groups.
- At 24 months, total body weight loss ranged from 12.2% in smokers to 14.9% in ex-smokers, with no statistically significant differences between groups.
- Follow-up adherence rates were comparable across groups, ranging from 73.5% to 93.1% at various time points.
- This material is relevant for clinicians counseling class 1 and 2 obese patients regarding endoscopic sleeve gastroplasty outcomes in relation to smoking habits.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflīx.eu/search/d/1b879234-743c-11ee-8c1a-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.