

VALIDATION OF MRI-MORPHOLOGIC RADIOLOGICAL DIAGNOSTIC CRITERIA IN A RETROSPECTIVE COHORT OF AUTOIMMUNE PANCREATITIS (AIP) PATIENTS

Max Seidensticker

UEG Week Copenhagen 2023

Poster

Pancreas

October 15, 2023

A retrospective blinded study of 156 MRI examinations evaluated five MR-morphologic criteria for distinguishing autoimmune pancreatitis from chronic pancreatitis, pancreatic cancer, primary sclerosing cholangitis, and non-HPB controls.

KEY POINTS

- Sausage-like enlargement showed the highest specificity for AIP (98.98% and 98.00% for two readers) but low sensitivity (30.91% and 26.79%).
- Restricted diffusion demonstrated moderate sensitivity (47.83% and 56.86%) and specificity (89.29% and 73.26%) with accuracy around 72-75% for both readers.
- Delayed enhancement in venous phase, icicle sign, and hyperenhancement of ductal wall all showed high specificity (79-100%) but low sensitivity (8-38%).
- Apparent diffusion coefficient values differed significantly between AIP, pancreatic cancer, and chronic pancreatitis, with AUROC of 0.85 for chronic pancreatitis and 0.68 for pancreatic cancer.
- Inter-reader agreement was moderate to substantial (kappa 0.23-0.85); this study is relevant for radiologists and gastroenterologists seeking to optimize MRI-based diagnosis of autoimmune pancreatitis.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/6ee59ce6-743c-11ee-a432-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.