gutflix
GI learning, personalised

Summary

Macrophages and metabolic liver disease

Moritz Peiseler

UEG Week Barcelona 2026 | Presentation

Hepatobiliary | Immunology | Digestive Oncology | Colorectal

October 18, 2026

Kupffer cells function as essential blood filters requiring specialized receptors and blood flow, but chronic liver injury causes macrophage heterogeneity and incomplete immunological recovery during fibrosis regression, with regression-associated macrophages driving exacerbated injury upon rechallenge.

KEY POINTS

- The speaker presented data showing kupffer cells express CRIG complement receptors and require adequate blood flow to catch and digest bacteria; mice lacking kupffer cells die rapidly from infection.
- In fibrotic livers, vascular remodeling creates large vessels where bacteria escape clearance, and most kupffer cells lose CRIG expression, impairing their bacteria-catching function.
- The speaker reported that in a Western diet MASLD model with minimal fibrosis, kupffer cells normalized completely after 24-week regression, whereas a high-fat diet model with robust fibrosis retained residual macrophages that could not catch bacteria.
- Using a CCl4 injury-regression-rechallenge model, the speaker identified regression-associated macrophages (RAMs) expressing high TREM2 that persist in former injury zones and trap neutrophils.
- Intravital microscopy showed neutrophils become immobilized on monocyte-derived macrophages during regression, and depletion of either macrophages or neutrophils before rechallenge reduced the exacerbated liver injury the speaker observed.
- The speaker stated that recent multi-omic sequencing suggests both population turnover and epigenetic changes contribute to macrophage memory, though hepatocytes from regression were more resistant to injury in culture experiments.

VIEW THIS CONTENT ON GUTFLIX

https://gutflix.eu/search/d/fc5f18b8-d14f-11f1-a491-daaa6dbee0a2

AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.