

VEGF-RELATED ANTIBODIOME CHARACTERIZES PATIENTS WITH CROHN'S DISEASE AND DIFFERENTIATES THE LOCATION OF INFLAMMATION

Piotr Eder

UEG Week Copenhagen 2023

Poster

IBD

October 15, 2023

A study of 66 patients with active Crohn's disease found that serum levels of autoantibodies against VEGF-A and its receptors distinguish patients with isolated colonic disease from those with ileal involvement.

KEY POINTS

- Compared to 197 healthy controls, Crohn's disease patients had significantly lower levels of autoantibodies against VEGF-A, VEGFR1, and VEGFR2, and significantly higher levels of autoantibodies against AT1R, with no difference in anti-ETAR levels.
- All VEGF-related autoantibodies were significantly lower in patients with isolated colonic disease compared to those with ileal involvement, discriminating between these locations with AUC values ranging from 0.72 to 0.77.
- In isolated colonic disease, correlations between autoantibodies were strongly positive ($r=0.5-0.9$) and CRP correlated positively with VEGF-A ($r=0.6$) but negatively with VEGF-related autoantibodies; these correlations were absent or less pronounced in patients with ileal involvement.
- The authors suggest this antibodyome profile may represent a laboratory signature for disease location in Crohn's disease, though these differences were independent of steroid and azathioprine treatment requirements.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/71872ea6-743c-11ee-b8a5-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.