

THE EFFECT OF S-ADEMETHIONINE ON PLASMA CITRULLINE LEVEL IN PATIENTS WITH CHRONIC LYMPHOPROLIFERATIVE DISORDERS WITH CHEMOTHERAPY-INDUCED DIARRHEA

Roman Skrypnyk

UEG Week Copenhagen 2023

Poster

Colorectal

October 15, 2023

S-adenosylmethionine as adjuvant treatment attenuated chemotherapy-induced gastrointestinal injury in patients with chronic lymphoproliferative disorders, as indicated by higher plasma citrulline levels.

KEY POINTS

- In 24 patients with chronic lymphoproliferative disorders receiving chemotherapy regimens (BR, FCR, or R-CHOP), S-adenosylmethionine (1000 mg/day IV for 10 days, then 500 mg twice daily for 20 days) was compared to chemotherapy alone over 3 courses.
- Chemotherapy alone was associated with enterocyte injury, demonstrated by 1.66 times lower plasma citrulline levels ($p=0.0002$) compared to baseline in the chemotherapy-only group.
- S-adenosylmethionine treatment resulted in 1.23 times higher blood citrulline levels ($p=0.0005$) after chemotherapy compared to the chemotherapy-only group, suggesting attenuation of intestinal dysfunction.
- Patients with chronic lymphoproliferative disorders had pre-existing mucosal injury before chemotherapy, with 1.25 to 1.26 times higher blood NANA concentrations than healthy controls.
- This material is most relevant for oncologists and gastroenterologists managing gastrointestinal complications in patients receiving chemotherapy for chronic lymphoproliferative disorders.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/145cd2fc-743d-11ee-8063-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.