

SPLENIC STIFFNESS IN NON-ALCOHOLIC FATTY LIVER DISEASE: CORRELATION TO SERUM BIOMARKERS AND LIVER ULTRASOUND TECHNOLOGY OF FIBROSIS

Rym Ennaifer

UEG Week Copenhagen 2023

Poster

Hepatobiliary

October 15, 2023

A prospective study of 44 NAFLD patients found splenic elasticity measurement significantly associated with liver elasticity and predictive of advanced fibrosis, with a threshold of 17.4 Kpa showing 85.7% specificity and 73.5% sensitivity.

KEY POINTS

- Splenic elasticity was significantly associated with liver elasticity ($p=0.012$) with an AUROC of 0.784, and a 17.4 Kpa threshold predicted advanced fibrosis with 85.7% specificity and 73.5% sensitivity.
- Among 44 patients with hepatic steatosis, Fibroscan-assessed fibrosis staging showed F0-F1 in 31 cases, F2 in 5 cases, F3 in 5 cases, and F4 in 3 cases.
- Advanced fibrosis correlated with NFS score ($p=0.016$) and FIB-4 ($p=0.001$), but splenic elasticity showed no association with NFS or FIB-4 scores.
- The authors concluded that splenic elasticity measurement is a simple and reliable non-invasive tool, comparable to liver elasticity, for predicting liver fibrosis in NAFLD.
- This prospective study conducted July 2022 to January 2023 may interest hepatologists evaluating non-invasive fibrosis assessment methods in NAFLD populations.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/223975d8-743d-11ee-a2de-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.