

ASSOCIATION BETWEEN HELICOBACTER PYLORI INFECTION AND GASTROESOPHAGEAL REFLUX DISEASE: MENDELIAN RANDOMIZATION STUDY

Shao-Hua Xie

UEG Week Copenhagen 2023

Poster

Stomach & H. Pylori

October 15, 2023

A Mendelian randomization study using data from FinnGen and UK Biobank found no evidence for a causal relation between *Helicobacter pylori* infection and the risk of gastroesophageal reflux disease.

KEY POINTS

- Eighteen single nucleotide polymorphisms associated with *Helicobacter pylori* IgG and six specific antibodies (UreA, CagA, VacA, Catalase, OMP, and GroEL) were used as genetic instruments to examine causal associations with GERD risk.
- Genetically predicted *Helicobacter pylori* IgG antibody levels were not associated with GERD risk in either FinnGen (OR=0.992, 95%CI 0.961-1.025) or UK Biobank (OR=0.999, 95%CI 0.998-1.001).
- Genetically predicted levels of the six *Helicobacter pylori* specific antibodies were not associated with GERD risk, except for UreA antibody in the MR-Egger method only, for which pleiotropy was indicated.
- Multivariable Mendelian randomization analysis also found no associations between *Helicobacter pylori* specific antibody levels and GERD risk.
- This work addresses inconsistent findings from previous epidemiological studies and may interest researchers and clinicians evaluating the role of *Helicobacter pylori* in GERD pathogenesis.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/934cb30e-743b-11ee-b2e9-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.