

VOLUMETRIC MEASUREMENT OF CROHN'S DISEASE ON MAGNETIC RESONANCE ENTEROGRAPHY: FEASIBILITY, CLINICAL UTILITY, AND ROLE IN ASSESSING TREATMENT RESPONSE

Stuart Taylor

UEG Week Copenhagen 2023

Poster

IBD

October 15, 2023

Volumetric measurement of terminal ileal Crohn's disease on MRE is feasible, correlates with endoscopic disease activity, and detects treatment response to biologics.

KEY POINTS

- In 30 patients, median disease volume was 20.9 cm³ in those with endoscopically active terminal ileal CD (CDEIS ≥3) compared to 5.7 cm³ in those without active disease on endoscopy.
- In 6 clinically responding patients treated with biologics, median disease volume decreased from 28.5 cm³ pre-treatment to 11 cm³ post-treatment, representing a median percentage difference of -110%.
- Manual segmentation by two board-certified abdominal radiologists showed a mean difference of -3.0 cm³ with limits of agreement of -21.8 to 15.9 cm³.
- The authors conclude that volumetric measurement represents a novel, objective MRE biomarker for assessing disease activity and treatment response in CD, worthy of further validation.
- This work addresses limitations of current MRE activity scores such as sMARIA, which often fail to capture early treatment response and rely on 2D assessment rather than full disease volume.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/fd861bf6-743c-11ee-87fd-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.