

## THE PREDICTIVE VALUE OF IRON DEFICIENCY ANAEMIA AND THE SEVERITY OF ANAEMIA ON THE ONE-YEAR DISEASE COURSE OF INFLAMMATORY BOWEL DISEASE

Tamás Molnár

UEG Week Copenhagen 2023

Poster

IBD

October 15, 2023

**In a retrospective cohort of 713 IBD patients followed for one year, iron deficiency anaemia without elevated inflammatory markers predicted worse outcomes including treatment escalation and major complications.**

### KEY POINTS

- Iron deficiency anaemia increased the risk of all complications (OR 2.745,  $p < 0.001$ ) and major complications including hospitalization (OR 2.606,  $p = 0.002$ ) but not abdominal surgery
- The severity of anaemia correlated with the prevalence of major complications ( $p < 0.001$ ), and CRP level was associated with anaemia severity ( $p < 0.001$ )
- Iron deficiency anaemia was defined as CRP  $> 10 \mu\text{g/ml}$ , ferritin  $< 30 \mu\text{g/ml}$ , transferrin saturation  $< 20\%$  with anaemia present; severe anaemia as haemoglobin  $< 100 \text{ g/l}$
- The study included 713 consecutive patients (median age 43 years, 402 Crohn's disease, 311 ulcerative colitis) from two centres with median 1.06 year follow-up
- The authors suggest that iron deficiency anaemia should be incorporated into treat-to-target strategies and monitored closely, with therapeutic intervention beyond iron replacement considered

VIEW THIS CONTENT ON GUTFLIX

<https://gutflrx.eu/search/d/0d42298c-743c-11ee-a1a5-0242ac140004>



**AI-generated summary**

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.