

TECHNICAL DIFFICULTY AND CLINICAL COURSES OF ENDOSCOPIC HAND SUTURING AFTER GASTRIC ENDOSCOPIC SUBMUCOSAL DISSECTION WITH SPECIAL REFERENCE TO THE LOCATION

Tsugumi Habu

UEG Week Copenhagen 2023

Poster

Oesophagus

October 15, 2023

Endoscopic hand suturing for mucosal defect closure after gastric endoscopic submucosal dissection achieved 99% complete closure but showed location-dependent technical difficulty and dehiscence rates in 74 lesions.

KEY POINTS

- Technical success was lower in the upper stomach (90%) compared to middle and lower regions (both 100%, $p=0.039$), while success rates did not differ significantly by wall location
- Closure was maintained at second-look endoscopy in 88% overall, with a trend toward lower maintenance on the lesser curvature (75%) compared to other walls (93-100%, $p=0.062$)
- Delayed bleeding occurred in 3% of cases with no severe adverse events, despite 68% of patients continuing antithrombotic agents perioperatively
- Mean suturing time was 41 minutes with 7 stitches (5.5 min/stitch) for defects averaging 33 mm diameter, with no significant speed differences by location
- The study suggests EHS on the upper stomach requires greater operator experience and that lesser curvature lesions need more careful technique to reduce dehiscence risk

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/46ef3658-743b-11ee-88b1-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.