

GENDER-SPECIFIC DIFFERENCES IN GASTROINTESTINAL SYMPTOMS AMONG PEDIATRIC PATIENTS WITH INFLAMMATORY BOWEL DISEASES AND COELIAC DISEASES ASSESSED WITH THE VALIDATED PSAGIS SCALE

Umair Anwar

UEG Week Copenhagen 2023

Poster

IBD

October 15, 2023

A study using the pSAGIS patient-reported outcome measure in 1,559 paediatric patients found significant gender differences in gastrointestinal symptom burden in Crohn's disease and ulcerative colitis but not in coeliac disease.

KEY POINTS

- In Crohn's disease (n=430), female patients had significantly higher overall GI symptom scores than males (7.3 ± 10.1 vs 2.8 ± 4.2 , $p < 0.001$), with higher scores across all five symptom groups including abdominal pain, dyspepsia, diarrhea, constipation, and dysphagia/nausea.
- In ulcerative colitis (n=302), male patients had significantly higher overall symptom scores than females (5.6 ± 8.5 vs 2.9 ± 5.9 , $p = 0.002$), driven particularly by higher dyspepsia and diarrhea scores in males.
- In coeliac disease (n=827), overall GI symptom scores were low and no significant gender differences were observed (3.1 ± 4.6 in males vs 3.9 ± 6.4 in females, $p = 0.064$).
- The authors conclude that these gender-specific symptom patterns may affect time to diagnosis, treatment initiation, and adherence, warranting further investigation.
- This study used the pSAGIS scale, a validated 21-item patient-reported measure assessing type and severity of GI symptoms, in a paediatric outpatient clinic in Vienna between 2016 and 2020.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/287e0ea4-743d-11ee-bbbe-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.