

QUANTITATIVE FLUORESCENCE MOLECULAR ENDOSCOPY USING CETUXIMAB-800CW TO EVALUATE RESPONSE TO NEOADJUVANT CHEMORADIOOTHERAPY IN LOCALLY ADVANCED RECTAL CANCER; LESSONS LEARNED

Wouter B. Nagengast

UEG Week Copenhagen 2023

Poster

Colorectal

October 15, 2023

Cetuximab-800CW fluorescence molecular endoscopy could not distinguish pathological complete response from residual tumor in rectal cancer patients after neoadjuvant chemoradiotherapy due to submucosal tumor localization and off-target macrophage binding.

KEY POINTS

- In a study of ten locally advanced rectal cancer patients after neoadjuvant chemoradiotherapy, five achieved pathological complete response and five had partial response.
- Quantitative fluorescence molecular endoscopy using cetuximab-800CW showed significantly higher fluorescent signal in former tumor areas compared to healthy tissue in both complete responders ($p < 0.0001$) and partial responders ($p < 0.0001$), but could not distinguish between the two groups.
- Fluorescence microscopy revealed specific binding of cetuximab-800CW to tumor epithelium occurred only in the submucosa, not at the luminal surface accessible to endoscopy.
- Aspecific binding to mucosal macrophages was observed in both complete and partial responders, resulting in enhanced off-target fluorescence signals that confounded interpretation.
- The authors suggest novel techniques measuring fluorescence below the superficial layer, such as ultrasound-guided needle biopsy with single fiber fluorescence spectroscopy, may overcome these limitations.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/3833611a-743c-11ee-ab36-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.