

FEASIBILITY AND SAFETY OF SPYGLASS DS CHOLEDOCHOSCOPE AND ENDOSCOPIC RADIOFREQUENCY ABLATION IN THE DIAGNOSIS AND TREATMENT OF UNRESECTABLE EXTRAHEPATIC CHOLANGIOCARCINOMA

Xiaofeng Zhang

UEG Week Copenhagen 2023

Poster

Hepatobiliary

October 15, 2023

A retrospective study of 90 patients with extrahepatic bile duct obstruction found that combining Spyglass DS choledochoscopy with radiofrequency ablation in a single session for unresectable extrahepatic cholangiocarcinoma improved diagnostic sensitivity and reduced procedure numbers without increasing adverse events.

KEY POINTS

- Spyglass DS choledochoscopy biopsy achieved 87.5% diagnostic sensitivity for extrahepatic cholangiocarcinoma compared to 38.9% for cell brush biopsy, a statistically significant difference.
- Performing Spyglass DS choledochoscopy and radiofrequency ablation in the same session reduced the mean number of ERCP procedures from 2.59 to 1.00 compared to conventional sequential approach.
- All patients in both groups successfully completed endoscopic radiofrequency ablation with 100% procedure success rate.
- Postoperative adverse event rates were similar between simultaneous and conventional groups at 14.7% and 14.3% respectively, with no significant difference, and bilirubin improvement was comparable between groups.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/31f63538-743d-11ee-92e2-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.