

ASEPTIC ABSCESS SYNDROME (AAS) IN A PATIENT WITH DIVERSION COLITIS

Yehuda Ringel

UEG Week Copenhagen 2023

Poster

Colorectal

October 15, 2023

This case report describes a 66-year-old woman with aseptic abscess syndrome presenting as liver lesions that resolved after treatment of concurrent diversion colitis, suggesting a potential new association between these conditions.

KEY POINTS

- A patient with permanent colostomy following Hartman's procedure developed rectal bleeding, tenesmus, weight loss, anemia, and elevated cholestatic liver enzymes two years post-surgery.
- Sigmoidoscopy revealed active colitis in the rectal stump, and imaging identified new liver lesions with biopsies showing sterile abscesses (acute and chronic inflammation with micro abscess formation, negative for malignancy and infection).
- The diagnosis of aseptic abscess syndrome was suggested, possibly secondary to diversion colitis.
- Treatment with topical 5-ASA enemas resulted in resolution of the colitis, normalization of liver enzymes, and complete resolution of liver abscesses.
- The authors propose this as the first reported case of aseptic abscess syndrome associated with diversion colitis, with the resolution of liver disease following colitis treatment suggesting the conditions are related.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflrx.eu/search/d/79f81f78-743c-11ee-a765-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.