

MUCOSA-ASSOCIATED MICROBIOTA WITH RELATED METABOLITES IN PATIENTS WITH DUODENAL NEOPLASMS

Yoshito Itoh

UEG Week Copenhagen 2023

Poster

Oesophagus

October 15, 2023

A study of early duodenal cancer patients identified specific mucosa-associated microbiota and metabolite differences between duodenal tumor tissue and healthy controls that may be associated with tumor development.

KEY POINTS

- Cutibacterium, Faecalibacterium, and Brevundimonas were significantly higher in duodenal tumor surrounding mucosa, while Neisseria and Porphyromonas were significantly higher in healthy control mucosa
- Eight metabolites differed significantly between tumor and control tissue, with citrulline, choline, and NAD⁺ decreased in intramucosal carcinoma compared to healthy controls
- Principal component analysis revealed variable choline metabolism and methionine circuits in duodenal tumor tissue
- The study enrolled early duodenal cancer patients from April 2020 to October 2021 and used 16S rRNA gene sequencing and capillary electrophoresis-Fourier transform mass spectrometry to analyze mucus samples from tumor tissue, surrounding mucosa, and healthy controls

VIEW THIS CONTENT ON GUTFLIX

<https://gutflifx.eu/search/d/0c3aa35c-743c-11ee-88e5-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.