

TIME TRENDS OF ENVIRONMENTAL AND SOCIOECONOMIC RISK FACTORS IN PATIENTS WITH INFLAMMATORY BOWEL DISEASE OVER 40 YEARS OF DIFFERENT THERAPEUTIC ERAS: A POPULATION-BASED COHORT BETWEEN 1977-2020

Zsuzsanna Erdelyi

UEG Week Copenhagen 2023

Poster

IBD

October 15, 2023

A four-decade population-based study from Hungary found declining smoking rates in Crohn's disease patients but no other major changes in environmental or socioeconomic factors that would explain the rising IBD incidence.

KEY POINTS

- Analysis of 2,240 incident IBD patients (61.2% ulcerative colitis) from Veszprem, Hungary, diagnosed between 1977 and 2020, divided into three therapeutic era cohorts.
- Active smoking rates in Crohn's disease patients decreased significantly over time from 60.2% to 38.6% across the three cohorts, while ulcerative colitis smoking rates remained stable and low at approximately 15%.
- Appendectomy before diagnosis decreased significantly in both ulcerative colitis (from 6.4% to 2.3%) and Crohn's disease (from 27.3% to 6.9%) over the study period.
- No significant changes were observed in family history of IBD, urban versus rural living patterns, or most socioeconomic indicators across the four decades.
- The authors conclude that trends in known environmental factors do not explain the sharp increase in IBD incidence observed over recent decades.

VIEW THIS CONTENT ON GUTFLIX

<https://gutflix.eu/search/d/353fe5ce-743b-11ee-9a9d-0242ac140004>



AI-generated summary

This summary was generated by an AI large language model based on the content transcript. It is for informational purposes only and should not be considered a substitute for clinical judgment. Always rely on your professional expertise and the full clinical context when making clinical decisions.